

## Cosmological Constraints from Multiple Probes in the Dark Energy Survey

T. M. C. Abbott,<sup>1</sup> A. Alarcon,<sup>2,3</sup> S. Allam,<sup>4</sup> P. Andersen,<sup>5,6</sup> F. Andrade-Oliveira,<sup>7,8</sup> J. Annis,<sup>4</sup> J. Asorey,<sup>9</sup> A. Avelino,<sup>10</sup> S. Avila,<sup>11</sup> D. Bacon,<sup>11</sup> N. Banik,<sup>4</sup> B. A. Bassett,<sup>12,13</sup> E. Baxter,<sup>14</sup> K. Bechtol,<sup>15</sup> M. R. Becker,<sup>16</sup> G. M. Bernstein,<sup>14</sup> E. Bertin,<sup>17,18</sup> J. Blazek,<sup>19,20</sup> S. L. Bridle,<sup>21</sup> D. Brooks,<sup>22</sup> D. Brout,<sup>14</sup> D. L. Burke,<sup>23,24</sup> J. Calcino,<sup>5</sup> H. Camacho,<sup>25,8</sup> A. Campos,<sup>26,7</sup> A. Carnero Rosell,<sup>27,8</sup> D. Carollo,<sup>28</sup> M. Carrasco Kind,<sup>29,30</sup> J. Carretero,<sup>31</sup> F. J. Castander,<sup>2,3</sup> R. Cawthon,<sup>32</sup> P. Challis,<sup>10</sup> K. C. Chan,<sup>2,3</sup> C. Chang,<sup>33,34</sup> M. Childress,<sup>35</sup> A. Clocchiatti,<sup>36</sup> M. Crocce,<sup>2,3</sup> C. E. Cunha,<sup>23</sup> C. B. D'Andrea,<sup>14</sup> L. N. da Costa,<sup>8,37</sup> C. Davis,<sup>23</sup> T. M. Davis,<sup>5</sup> J. De Vicente,<sup>27</sup> D. L. DePoy,<sup>38</sup> J. DeRose,<sup>39,23</sup> S. Desai,<sup>40</sup> H. T. Diehl,<sup>4</sup> J. P. Dietrich,<sup>41,42</sup> S. Dodelson,<sup>26</sup> P. Doel,<sup>22</sup> A. Drlica-Wagner,<sup>4,34</sup> T. F. Eifler,<sup>43,44</sup> J. Elvin-Poole,<sup>19,45</sup> J. Estrada,<sup>4</sup> A. E. Evrard,<sup>46,47</sup> E. Fernandez,<sup>31</sup> A. V. Filippenko,<sup>48,49</sup> B. Flaugher,<sup>4</sup> R. J. Foley,<sup>50</sup> P. Fosalba,<sup>2,3</sup> J. Frieman,<sup>4,34</sup> L. Galbany,<sup>51</sup> J. García-Bellido,<sup>52</sup> M. Gatti,<sup>31</sup> E. Gaztanaga,<sup>2,3</sup> D. W. Gerdes,<sup>46,47</sup> T. Giannantonio,<sup>53,54,55</sup> K. Glazebrook,<sup>56</sup> D. A. Goldstein,<sup>57</sup> D. Gruen,<sup>23,24</sup> R. A. Gruendl,<sup>29,30</sup> J. Gschwend,<sup>8,37</sup> G. Gutierrez,<sup>4</sup> W. G. Hartley,<sup>22,58</sup> S. R. Hinton,<sup>5</sup> D. L. Hollowood,<sup>50</sup> K. Honscheid,<sup>19,45</sup> J. K. Hoormann,<sup>5</sup> B. Hoyle,<sup>59,55</sup> D. Huterer,<sup>47</sup> B. Jain,<sup>14</sup> D. J. James,<sup>60</sup> M. Jarvis,<sup>14</sup> T. Jeltema,<sup>50</sup> E. Kasai,<sup>61,13</sup> S. Kent,<sup>4,34</sup> R. Kessler,<sup>33,34</sup> A. G. Kim,<sup>62</sup> R. P. Kirshner,<sup>63,64</sup> N. Kokron,<sup>39,23</sup> E. Krause,<sup>43</sup> R. Kron,<sup>4,34</sup> K. Kuehn,<sup>65</sup> N. Kuropatkin,<sup>4</sup> O. Lahav,<sup>22</sup> J. Lasker,<sup>33,34</sup> P. Lemos,<sup>22,53,54</sup> G. F. Lewis,<sup>66</sup> T. S. Li,<sup>4,34</sup> C. Lidman,<sup>67</sup> M. Lima,<sup>25,8</sup> H. Lin,<sup>4</sup> E. Macaulay,<sup>11</sup> N. MacCrann,<sup>19,45</sup> M. A. G. Maia,<sup>8,37</sup> K. S. Mandel,<sup>68</sup> M. March,<sup>14</sup> J. Marriner,<sup>4</sup> J. L. Marshall,<sup>38</sup> P. Martini,<sup>19,69</sup> R. G. McMahon,<sup>53,54</sup> P. Melchior,<sup>70</sup> F. Menanteau,<sup>29,30</sup> R. Miquel,<sup>71,31</sup> J. J. Mohr,<sup>41,42,59</sup> E. Morganson,<sup>30</sup> J. Muir,<sup>23</sup> A. Möller,<sup>72,67</sup> E. Neilsen,<sup>4</sup> R. C. Nichol,<sup>11</sup> B. Nord,<sup>4</sup> R. L. C. Ogando,<sup>8,37</sup> A. Palmese,<sup>4</sup> Y.-C. Pan,<sup>73,74</sup> H. V. Peiris,<sup>22</sup> W. J. Percival,<sup>75,76</sup> A. A. Plazas,<sup>44</sup> A. Porredon,<sup>2,3</sup> J. Prat,<sup>31</sup> A. K. Romer,<sup>77</sup> A. Roodman,<sup>23,24</sup> R. Rosenfeld,<sup>78,8</sup> A. J. Ross,<sup>19</sup> E. S. Rykoff,<sup>23,24</sup> S. Samuroff,<sup>26</sup> C. Sánchez,<sup>14</sup> E. Sanchez,<sup>27</sup> V. Scarpine,<sup>4</sup> R. Schindler,<sup>24</sup> M. Schubnell,<sup>47</sup> D. Scolnic,<sup>34</sup> L. F. Secco,<sup>14</sup> S. Serrano,<sup>2,3</sup> I. Sevilla-Noarbe,<sup>27</sup> R. Sharp,<sup>67</sup> E. Sheldon,<sup>79</sup> M. Smith,<sup>35</sup> M. Soares-Santos,<sup>80</sup> F. Sobreira,<sup>81,8</sup> N. E. Sommer,<sup>72,67</sup> E. Swann,<sup>11</sup> M. E. C. Swanson,<sup>30</sup> G. Tarle,<sup>47</sup> D. Thomas,<sup>11</sup> R. C. Thomas,<sup>62</sup> M. A. Troxel,<sup>19,45</sup> B. E. Tucker,<sup>72,67</sup> S. A. Uddin,<sup>82</sup> P. Vielzeuf,<sup>31</sup> A. R. Walker,<sup>1</sup> M. Wang,<sup>4</sup> N. Weaverdyck,<sup>47</sup> R. H. Wechsler,<sup>39,23,24</sup> J. Weller,<sup>41,59,55</sup> B. Yanny,<sup>4</sup> B. Zhang,<sup>72,67</sup> Y. Zhang,<sup>4</sup> and J. Zuntz<sup>83</sup>

(DES Collaboration)

<sup>1</sup>*Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatory, Casilla 603, La Serena, Chile*

<sup>2</sup>*Institut d'Estudis Espacials de Catalunya (IEEC), 08034 Barcelona, Spain*

<sup>3</sup>*Institute of Space Sciences (ICE, CSIC), Campus UAB, Carrer de Can Magrans, s/n, 08193 Barcelona, Spain*

<sup>4</sup>*Fermi National Accelerator Laboratory, P. O. Box 500, Batavia, IL 60510, USA*

<sup>5</sup>*School of Mathematics and Physics, University of Queensland, Brisbane, QLD 4072, Australia*

<sup>6</sup>*University of Copenhagen, Dark Cosmology Centre, Juliane Maries Vej 30, 2100 Copenhagen O*

<sup>7</sup>*Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil*

<sup>8</sup>*Laboratório Interinstitucional de e-Astronomia - LInEA, Rua Gal. José Cristino 77, Rio de Janeiro, RJ - 20921-400, Brazil*

<sup>9</sup>*Korea Astronomy and Space Science Institute, Yuseong-gu, Daejeon, 305-348, Korea*

<sup>10</sup>*Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138, USA*

<sup>11</sup>*Institute of Cosmology and Gravitation, University of Portsmouth, Portsmouth, PO1 3FX, UK*

<sup>12</sup>*African Institute for Mathematical Sciences, 6 Melrose Road, Muizenberg, 7945, South Africa*

<sup>13</sup>*South African Astronomical Observatory, P.O.Box 9, Observatory 7935, South Africa*

<sup>14</sup>*Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104, USA*

<sup>15</sup>*LSST, 933 North Cherry Avenue, Tucson, AZ 85721, USA*

<sup>16</sup>*Argonne National Laboratory, 9700 South Cass Avenue, Lemont, IL 60439, USA*

<sup>17</sup>*CNRS, UMR 7095, Institut d'Astrophysique de Paris, F-75014, Paris, France*

<sup>18</sup>*Sorbonne Universités, UPMC Univ Paris 06, UMR 7095, Institut d'Astrophysique de Paris, F-75014, Paris, France*

<sup>19</sup>*Center for Cosmology and Astro-Particle Physics, The Ohio State University, Columbus, OH 43210, USA*

<sup>20</sup>*Institute of Physics, Laboratory of Astrophysics, École Polytechnique Fédérale de Lausanne (EPFL), Observatoire de Sauverny, 1290 Versoix, Switzerland*

<sup>21</sup>*Jodrell Bank Center for Astrophysics, School of Physics and Astronomy, University of Manchester, Oxford Road, Manchester, M13 9PL, UK*

<sup>22</sup>*Department of Physics & Astronomy, University College London, Gower Street, London, WC1E 6BT, UK*

<sup>23</sup>*Kavli Institute for Particle Astrophysics & Cosmology, P. O. Box 2450, Stanford University, Stanford, CA 94305, USA*

<sup>24</sup>*SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA*

<sup>25</sup>*Departamento de Física Matemática, Instituto de Física, Universidade de São Paulo, CP 66318, São Paulo, SP, 05314-970, Brazil*

<sup>26</sup>*Department of Physics, Carnegie Mellon University, Pittsburgh, Pennsylvania 15312, USA*

- <sup>27</sup> *Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain*
- <sup>28</sup> *INAF, Astrophysical Observatory of Turin, I-10025 Pino Torinese, Italy*
- <sup>29</sup> *Department of Astronomy, University of Illinois at Urbana-Champaign, 1002 W. Green Street, Urbana, IL 61801, USA*
- <sup>30</sup> *National Center for Supercomputing Applications, 1205 West Clark St., Urbana, IL 61801, USA*
- <sup>31</sup> *Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Campus UAB, 08193 Bellaterra (Barcelona) Spain*
- <sup>32</sup> *Physics Department, 2320 Chamberlin Hall, University of Wisconsin-Madison, 1150 University Avenue Madison, WI 53706-1390*
- <sup>33</sup> *Department of Astronomy and Astrophysics, University of Chicago, Chicago, IL 60637, USA*
- <sup>34</sup> *Kavli Institute for Cosmological Physics, University of Chicago, Chicago, IL 60637, USA*
- <sup>35</sup> *School of Physics and Astronomy, University of Southampton, Southampton, SO17 1BJ, UK*
- <sup>36</sup> *Millennium Institute of Astrophysics and Department of Physics and Astronomy, Universidad Católica de Chile, Santiago, Chile*
- <sup>37</sup> *Observatório Nacional, Rua Gal. José Cristino 77, Rio de Janeiro, RJ - 20921-400, Brazil*
- <sup>38</sup> *George P. and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, and Department of Physics and Astronomy, Texas A&M University, College Station, TX 77843, USA*
- <sup>39</sup> *Department of Physics, Stanford University, 382 Via Pueblo Mall, Stanford, CA 94305, USA*
- <sup>40</sup> *Department of Physics, IIT Hyderabad, Kandi, Telangana 502285, India*
- <sup>41</sup> *Excellence Cluster Universe, Boltzmannstr. 2, 85748 Garching, Germany*
- <sup>42</sup> *Faculty of Physics, Ludwig-Maximilians-Universität, Scheinerstr. 1, 81679 Munich, Germany*
- <sup>43</sup> *Department of Astronomy/Steward Observatory, 933 North Cherry Avenue, Tucson, AZ 85721-0065, USA*
- <sup>44</sup> *Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Dr., Pasadena, CA 91109, USA*
- <sup>45</sup> *Department of Physics, The Ohio State University, Columbus, OH 43210, USA*
- <sup>46</sup> *Department of Astronomy, University of Michigan, Ann Arbor, MI 48109, USA*
- <sup>47</sup> *Department of Physics, University of Michigan, Ann Arbor, MI 48109, USA*
- <sup>48</sup> *Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA*
- <sup>49</sup> *Miller Senior Fellow, Miller Institute for Basic Research in Science, University of California, Berkeley, CA 94720, USA*
- <sup>50</sup> *Santa Cruz Institute for Particle Physics, Santa Cruz, CA 95064, USA*
- <sup>51</sup> *PITT PACC, Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, PA 15260, USA*
- <sup>52</sup> *Instituto de Física Teórica UAM/CSIC, Universidad Autónoma de Madrid, 28049 Madrid, Spain*
- <sup>53</sup> *Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK*
- <sup>54</sup> *Kavli Institute for Cosmology, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK*
- <sup>55</sup> *Universitäts-Sternwarte, Fakultät für Physik, Ludwig-Maximilians Universität München, Scheinerstr. 1, 81679 München, Germany*
- <sup>56</sup> *Centre for Astrophysics & Supercomputing, Swinburne University of Technology, Victoria 3122, Australia*
- <sup>57</sup> *California Institute of Technology, 1200 East California Blvd, MC 249-17, Pasadena, CA 91125, USA*
- <sup>58</sup> *Department of Physics, ETH Zurich, Wolfgang-Pauli-Strasse 16, CH-8093 Zurich, Switzerland*
- <sup>59</sup> *Max Planck Institute for Extraterrestrial Physics, Giessenbachstrasse, 85748 Garching, Germany*
- <sup>60</sup> *Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA*
- <sup>61</sup> *Department of Physics, University of Namibia, 340 Mandume Ndemufayo Avenue, Pionierspark, Windhoek, Namibia*
- <sup>62</sup> *Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA*
- <sup>63</sup> *Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138, USA*
- <sup>64</sup> *Gordon and Betty Moore Foundation, 1661 Page Mill Road, Palo Alto, CA 94304, USA*
- <sup>65</sup> *Australian Astronomical Optics, Macquarie University, North Ryde, NSW 2113, Australia*
- <sup>66</sup> *Sydney Institute for Astronomy, School of Physics, A28, The University of Sydney, NSW 2006, Australia*
- <sup>67</sup> *The Research School of Astronomy and Astrophysics, Australian National University, ACT 2601, Australia*
- <sup>68</sup> *Institute of Astronomy and Kavli Institute for Cosmology, Madingley Road, Cambridge, CB3 0HA, UK*
- <sup>69</sup> *Department of Astronomy, The Ohio State University, Columbus, OH 43210, USA*
- <sup>70</sup> *Department of Astrophysical Sciences, Princeton University, Peyton Hall, Princeton, NJ 08544, USA*
- <sup>71</sup> *Institució Catalana de Recerca i Estudis Avançats, E-08010 Barcelona, Spain*
- <sup>72</sup> *ARC Centre of Excellence for All-sky Astrophysics (CAASTRO)*
- <sup>73</sup> *Division of Theoretical Astronomy, National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan*
- <sup>74</sup> *Institute of Astronomy and Astrophysics, Academia Sinica, Taipei 10617, Taiwan*
- <sup>75</sup> *Department of Physics and Astronomy, University of Waterloo, 200 University Ave W, Waterloo, ON N2L 3G1, Canada*
- <sup>76</sup> *Perimeter Institute for Theoretical Physics, 31 Caroline St. North, Waterloo, ON N2L 2Y5, Canada*
- <sup>77</sup> *Department of Physics and Astronomy, Pevensey Building, University of Sussex, Brighton, BN1 9QH, UK*
- <sup>78</sup> *ICTP South American Institute for Fundamental Research  
Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil*
- <sup>79</sup> *Brookhaven National Laboratory, Bldg 510, Upton, NY 11973, USA*
- <sup>80</sup> *Brandeis University, Physics Department, 415 South Street, Waltham MA 02453*
- <sup>81</sup> *Instituto de Física Gleb Wataghin, Universidade Estadual de Campinas, 13083-859, Campinas, SP, Brazil*
- <sup>82</sup> *Observatories of the Carnegie Institution for Science, 813 Santa Barbara St., Pasadena, CA 91101, USA*
- <sup>83</sup> *Institute for Astronomy, University of Edinburgh, Edinburgh EH9 3HJ, UK*

(Dated: November 8, 2018)

The combination of multiple observational probes has long been advocated as a powerful way to constrain cosmological parameters, in particular dark energy. The Dark Energy Survey has measured 207 spectroscopically-confirmed Type Ia supernova lightcurves; the baryon acoustic oscillation feature; weak gravitational lensing; and galaxy clustering. Here we present combined results from these probes, deriving constraints on the equation of state,  $w$ , of dark energy and its energy density in the Universe. Independently of other experiments, such as those that measure the cosmic microwave background, the probes from this single photometric survey rule out a Universe with no dark energy, finding  $w = -0.80^{+0.09}_{-0.11}$ . The geometry is shown to be consistent with a spatially flat Universe, and we report a constraint on the baryon density of  $\Omega_b = 0.069^{+0.009}_{-0.012}$  that is independent of early Universe measurements. These results demonstrate the potential power of large multi-probe photometric surveys and pave the way for order of magnitude advances in our constraints on properties of dark energy and cosmology over the next decade.

Keywords: dark energy; dark matter; cosmology: observations; cosmological parameters

## INTRODUCTION

The discovery of the accelerating Universe [1, 2] was perhaps the crowning achievement of 20th century cosmology, as it indicated the presence of a qualitatively new component in the Universe that dominates the expansion in the last several billion years. The nature of dark energy — the component that causes the accelerated expansion — is unknown, and understanding its properties and origin is one of the principal challenges in modern physics. Current measurements are consistent with an interpretation of dark energy as a cosmological constant in General Relativity. Any deviation from this interpretation in space or time would constitute a landmark discovery in fundamental physics [3].

Dark energy leaves imprints on cosmological observations, typically split into two regimes — 1) it modifies the geometry of the Universe, increasing distances and volumes in the Universe over time via the accelerated expansion, and 2) it suppresses the growth of cosmic structure. However, these effects can be mimicked by the variation of other cosmological parameters, including the dark matter density and curvature. Consequently, measuring dark energy properties requires a combination of cosmological probes that are sensitive to both classes of effects to break these parameter degeneracies [4–6].

Historically, the most powerful cosmic probe has been the cosmic microwave background (CMB) [7–9], relic radiation from the surface of last scattering only 400,000 years after the Big Bang. Low-redshift probes measure the Universe as it exists in the last several billion years, when dark energy dominates the expansion. Comparing or combining constraints between the CMB and lower redshift measurements requires us to extrapolate predictions for the present-day Universe starting from initial conditions over 13 billion years ago. This can be a powerful test of our models, but it requires precise, independent constraints from low-redshift experiments. Low-redshift probes include Type Ia supernova (SNe Ia) measurements, which treat the SNe Ia as standardizable candles and employ redshift and flux measurements to probe the redshift-luminosity distance relation [10]; baryon acoustic oscillations (BAO), which use a ‘standard ruler’ scale in the cosmic density field, imprinted by sound waves at recombination, to probe various redshift-distance combinations [11]; galaxy clustering, which measures the density field up

to some bias between galaxy density and the underlying dark matter density, and redshift-space distortions (RSD) in the clustering [12]; the counts of galaxy clusters, representing the most extreme density peaks in the Universe [13]; strong gravitational lensing [14]; and weak gravitational lensing, which probes changes in the gravitational potential along the line of sight using coherent distortions in observed properties of galaxies or the CMB, e.g. to measure the dark and baryonic matter distribution [15].

We report here the first results from the Dark Energy Survey (DES) combining precision probes of both geometry and growth of structure that include BAO, SNe Ia, and weak lensing and galaxy clustering. DES has previously shown separate cosmological constraints using weak lensing and galaxy clustering [16], BAO [17], and SNe Ia [18]. We now combine these probes and begin to fully realize the power of this multi-probe experiment to produce independent measurements of the properties of dark energy.

The work presented here demonstrates our ability to extract and combine diverse cosmological observables from large-scale surveys of the evolved Universe. Previous dark energy constraints have relied on combining the likelihoods of many separate and independent experiments to produce precise constraints on cosmological models including dark energy. For this traditional approach each experiment has performed an independent analysis to validate measurements and has separate calibration methodologies and requirements, thus ensuring that many potential systematics are uncorrelated between probes. The DES analysis presented here, however, uses a common set of both calibration methodologies and systematics modeling and marginalization across probes, which enables a consistently validated analysis. Perhaps most importantly, this common framework allows us to standardize requirements like blinding across these probes, which is essential to minimize the impact of experimenter bias [19].

The fundamental interest in understanding the nature of dark energy has spurred the development of multiple large photometric surveys that image the sky, capable of independently combining multiple cosmic probes. The current generation of surveys includes the Hyper-Suprime Cam survey (HSC) [20], the Kilo-Degree Survey (KiDS) [21], and the focus of this work, DES [22]. The next generation of these surveys will include the Large Synoptic Survey Telescope

(LSST) [23], a ground-based telescope that will observe the entire southern hemisphere with very high cadence, and space telescopes Euclid [24] and the Wide-Field InfraRed Survey Telescope (WFIRST) [25]. In parallel with imaging surveys, the distribution of galaxies measured by spectroscopic surveys (i.e., BOSS [26], eBOSS [27], and the planned DESI [28] and PFS [29] surveys) provides powerful constraints on the distance-redshift relation via BAO measurements and the growth of structure via redshift space distortions. The union of these results over the following several years, and into the next decades, will ensure that we are able to take advantage of the benefits of multiple independent, self-consistent, and blinded multi-probe analyses like we present here for DES.

## COSMIC PROBES

### The Dark Energy Survey

DES cosmic probes span a wide range of redshifts up to  $z \approx 1.3$ , and include weak gravitational lensing and galaxy clustering due to large-scale structure [16], SNe Ia [18], and BAO [17]. Each of these are able to constrain dark energy independently and their combination is even more powerful. These probes utilize a subset of data from DES taken during its first three observing seasons (Aug. 2013 to Feb. 2016). Spectroscopically confirmed SNe Ia are identified from images in all three seasons (DES Y3) in  $27 \text{ deg}^2$  of repeated deep-field observations, while weak lensing and large-scale structure information is derived from images taken only in the first season (DES Y1), ending Feb. 2014 and covering  $1321 \text{ deg}^2$  of the southern sky in *grizY* filters. DES uses the 570-megapixel Dark Energy Camera (DECam [30]) at the Cerro Tololo Inter-American Observatory (CTIO) 4m Blanco telescope in Chile. By the end of DES observations in January 2019, we anticipate an order of magnitude increase in the number of useable SNe, while the area of sky used for the other probes will increase by a factor of three to  $5000 \text{ deg}^2$ . Analysis of the later years of survey data is ongoing.

Data is processed through the DES Data Management system [31–34]. This system detrends and calibrates the raw images, creates coadded images from individual exposures, and detects and catalogs astrophysical objects. This catalog is further cleaned and calibrated to create a high-quality (‘Gold’) object catalog [35] from which weak lensing and large-scale structure measurements are made. The deep fields are also processed through a separate difference imaging pipeline to identify transients [36, 37]. The photometric and astrometric calibrations are common to all cosmology probes discussed below.

#### Weak Gravitational Lensing and Large-Scale Structure

For weak gravitational lensing measurements, we use the measured shapes and positions of 26 million galaxies in the

redshift range  $0.2 < z < 1.3$ , split into four redshift bins. The galaxy shapes are measured via the METACALIBRATION method [38, 39] using *riz*-band exposures [40]. Photometric redshifts for the objects are determined from a modified version of the BPZ method [41], described and calibrated in Ref. [42].

For measurements of the angular galaxy clustering, we utilize the positions of a sample of luminous red galaxies that have precise photometric redshifts selected with the REDMAGIC algorithm [43]. This results in a sample of 650,000 galaxies over the redshift range  $0.15 < z < 0.9$ , split into five narrow redshift bins. Residual correlations of number density with survey conditions in the REDMAGIC sample are calibrated in Ref. [44]. The precise redshifts of REDMAGIC galaxies allow us to infer information about the much larger photo- $z$  bias in the weak lensing catalog. The photo- $z$  calibration methodology is consistent between the weak lensing and REDMAGIC samples.

We use measurements from each of these galaxy samples to construct a set of three two-point correlation function observables we label ‘ $3 \times 2\text{pt}$ ’. These include the galaxy shear auto-correlation (cosmic shear), the galaxy position-shear cross-correlation (galaxy-galaxy lensing), and the galaxy position auto-correlation (galaxy clustering). The analysis was described in a series of papers that include the covariance and analysis framework [45, 46], the measurements and validation [44, 47–49], and the cosmological results [16]. We utilize the ‘ $3 \times 2\text{pt}$ ’ likelihood pipeline from this set of papers as implemented in COSMOSIS [50]. This combination of probes produces a tight constraint on the amplitude of matter clustering in the Universe and on the properties of dark energy over the last six billion years.

#### Type Ia Supernovae

The DES-SN sample is comprised of 207 spectroscopically confirmed SNe Ia in the redshift range  $0.07 < z < 0.85$ . The sample-building and analysis pipelines are discussed in a series of papers that detail the SN Ia search and discovery [34, 36, 37]; spectroscopic follow-up [51]; photometry [52]; calibration [53, 54]; simulations [55]; and technique of accounting for selection bias [56]. The analysis methodology and systematic uncertainties are presented in Ref. [57]. These results are used to constrain cosmology [18] and the Hubble constant [58]. In Refs. [18, 57, 58] the DES-SN sample is combined with a ‘Low- $z$ ’ ( $z < 0.1$ ) sample, which includes SNe from the Harvard-Smithsonian Center for Astrophysics surveys [59, 60] and the Carnegie Supernova Project [61]. Selection effects and calibration of these samples is discussed in [62]. Here we fit for DES-SN alone, and only include the Low- $z$  sample for comparison to Ref. [18]. We compute the SNe likelihood using the SNe module [62] implemented in COSMOSIS.

### Baryon Acoustic Oscillations

A sample of 1.3 million galaxies from the DES Y1 ‘Gold’ catalog in the redshift range  $0.6 < z < 1.0$  was used to measure the BAO scale in the distribution of galaxies. Details of the galaxy sample selection are in Ref. [63]. Calibrations of the galaxy selection function are consistently derived for the BAO and ‘3×2pt’ samples. This BAO measurement was presented in Ref. [17] and provides a likelihood for the ratio between the angular diameter distance to redshift 0.81,  $D_A(z = 0.81)$ , and the sound horizon at the drag epoch,  $r_d$ . This analysis used 1800 simulations [64] and three methods to compute the galaxy clustering [65–67]. The BAO likelihood is implemented in COSMOSIS. The galaxy samples used in the ‘3×2pt’ angular clustering measurements and in the BAO analysis share a common footprint in the sky and overlap significantly in volume over the redshift range  $0.6 < z < 0.9$ , which will produce some non-zero correlation between the two measurements. However, the intersection of the galaxy populations is only about 14% of the total BAO galaxy sample and we detect no significant BAO constraint when using the ‘3x2pt’ galaxy clustering measurements. We thus ignore this negligible correlation when combining the two probes.

#### External Data for Comparison

We use external constraints that combine state-of-the-art CMB, SNe Ia, and spectroscopic BAO measurements to compare our results against. For the CMB data, we utilize full-sky temperature ( $T$ ) and polarization ( $E$ - and  $B$ -mode) measurements from the Planck survey, combining  $TT$  ( $\ell \in [2, 2508]$ ), and  $EE$ ,  $BB$  and  $TE$  ( $\ell \in [2, 29]$ ) (commonly referred to as ‘TT+LowP’) [68] with weak lensing measurements derived from the temperature data [69]. We use the Planck likelihood from Ref. [70].

For external SNe Ia measurements, we use the Pantheon compilation [62]. Pantheon combines SNe Ia samples from Pan-STARRS1, SDSS, SNLS, various low- $z$  data sets, and HST. The Pantheon data set is based on the Pan-STARRS1 Supercal algorithm [71] that establishes a global calibration for the 13 different SNe Ia samples, with a total of 1048 SNe in  $0.01 < z < 2.26$ .

Finally, external spectroscopic BAO measurements are taken from BOSS DR12 [12], the 6DF Galaxy Survey [72], and the SDSS Main Galaxy Sample [73]. These measurements of the BAO scale span a redshift range of  $0.1 < z < 0.6$ .

#### CONSTRAINTS ON DARK ENERGY

We present here a dark energy analysis that combines for the first time the DES probes described above. DES is able to strongly constrain dark energy models without the CMB by probing over a wide redshift range ( $z \lesssim 1$ ) the growth of

TABLE I. Cosmological parameter constraints in the  $\Lambda$ CDM and  $w$ CDM models using only DES data. We report the 1D peak of the posterior and asymmetric 68% confidence limits. The marginalized parameters (and priors) are: the primordial perturbation amplitude  $10^9 A_s \in [0.5, 10.0]$ , the Hubble constant  $H_0 \in [55, 90]$  km s<sup>-1</sup>Mpc<sup>-1</sup>, the spectral index  $n_s \in [0.87, 1.07]$ , and the neutrino mass density  $\Omega_\nu h^2 \in [0.0006, 0.01]$ .

Parameter	$\Lambda$ CDM	$w$ CDM	Flat Prior
$\Omega_m$	$0.299^{+0.024}_{-0.020}$	$0.300^{+0.023}_{-0.021}$	[0.1, 0.9]
$\Omega_b$	$0.069^{+0.009}_{-0.012}$	$0.064^{+0.013}_{-0.009}$	[0.03, 0.12]
$\Omega_k$	$0.252^{+0.095}_{-0.14}$	0	[-0.1, 0.5]
$\Omega_\Lambda$	$0.47^{+0.14}_{-0.12}$	$0.700^{+0.021}_{-0.023}$	Derived
$w$	-1	$-0.80^{+0.09}_{-0.11}$	[-2, -0.33]
$S_8 \equiv \sigma_8 \sqrt{\Omega_m/0.3}$	$0.801^{+0.028}_{-0.026}$	$0.786^{+0.029}_{-0.019}$	Derived

structure and distance-redshift relation, which are both sensitive to the presence of dark energy. The dark energy equation of state  $w$  relates the pressure ( $P$ ) to the energy density ( $\rho$ ) of the dark energy fluid:  $w = P/\rho$ , where  $w = -1$  is equivalent to a cosmological constant  $\Lambda$  in the field equations. We probe the nature of dark energy in two ways: 1) we constrain the dark energy density relative to the critical density today,  $\Omega_\Lambda$ , assuming that dark energy takes the form of a cosmological constant and allowing non-zero curvature (the  $\Lambda$ CDM model), and 2) we measure  $w$  as a free parameter (the  $w$ CDM model) with fixed curvature ( $\Omega_k = 0$ ). The total energy density of the Universe today is composed of the sum of fractional components  $1 = \Omega_k + \Omega_m + \Omega_\Lambda$ , where the components are: curvature ( $\Omega_k$ ), the total matter ( $\Omega_m$ ), and dark energy ( $\Omega_\Lambda$ ). The radiation density is assumed to be negligible today. In both  $\Lambda$ CDM and  $w$ CDM models, we explore the ability of DES to constrain these properties of dark energy and compare this to the state-of-the-art constraints combining measurements from many external surveys. We follow the analysis methods and model definitions from Ref. [16], which includes varying the neutrino mass density in all models.

Figure 1 shows our constraints on  $\Omega_\Lambda$  in the  $\Lambda$ CDM model, where  $w = -1$ . We combine our ‘3×2pt’, SNe Ia (with-out the external Low- $z$  sample), and photometric BAO measurements to constrain  $\Omega_\Lambda$  and  $\Omega_m$ . This is compared to the constraint from the external data sets. Using DES data we are able to independently confirm the existence of a dark energy component in the Universe at  $4\sigma$  significance. This is the first time a photometric survey has independently made a significant constraint on the energy density of both dark energy and dark matter without assuming a flat model based on early Universe constraints. It represents an important milestone for future analyses from DES and surveys like Euclid, LSST, and WFIRST.

In Fig. 2, we show the constraint on  $w$  and  $\Omega_m$ , assuming the  $w$ CDM model. We show the same comparison with






FIG. 1. Constraints on the present-day dark energy density  $\Omega_\Lambda$  and matter density  $\Omega_m$ , relative to the critical density, in an  $\Lambda$ CDM model with free curvature and neutrino mass density. We compare the constraint from DES data alone (black contours), including information from weak gravitational lensing, large-scale structure, SNe Ia, and photometric BAO, to the best available external data (green contours), combining information from the CMB, SNe Ia, and spectroscopic BAO. We identify the flat model ( $\Omega_k = 0$ ) with a dotted line and distinguish accelerating and decelerating universes with a dashed line. Contours represent the 68% and 95% confidence limits (CL).

external data as in Fig. 1, but also include a case where we supplement DES-discovered SNe Ia with the Low- $z$  SNe sample to anchor the SNe redshift-distance relation at low redshift following Ref. [18]. This low-redshift SNe anchor is a common and significant contribution to both the DES+Low- $z$  and external constraints on  $w$ . In all cases, the existing data are consistent with a cosmological constant ( $w = -1$ ). This subset of the final DES data constrains  $w$  to within a factor of  $\sim 2$  of that obtained when combining multiple major external data sets. This clearly illustrates the prospects for multiple independent, precise low-redshift constraints on dark energy from upcoming large-scale photometric experiments.

The constraints on all model parameters are summarized in Table I. In the  $\Lambda$ CDM model, DES is also able to constrain the total matter density to 7% (68% CL), the baryon density to 15%, and the correlation amplitude to 3%, described by  $S_8 \equiv \sigma_8 \sqrt{\Omega_m}/0.3$ , where  $\sigma_8$  measures the current-day clustering amplitude. The constraints are comparable in  $w$ CDM. Fixing  $\Omega_k = 0$ , we find the  $S_8$  constraint is improved by a factor of 1.2, but there is otherwise no significant improvement in other parameters. These parameter constraints beyond dark energy are driven by the ‘3 $\times$ 2pt’ measurement. In particular,




FIG. 2. Constraints on the dark energy equation of state  $w$  and  $\Omega_m$  in a  $w$ CDM model with fixed curvature ( $\Omega_k = 0$ ) and free neutrino mass density. We compare constraints from the DES data alone (black contours) to the best available external data (green contours), as in Fig. 1, but also show the impact of including a low-redshift SNe Ia data set (Low- $z$ ) to anchor the DES SNe Ia as done in Ref. [18] (blue contours). Each component of the DES analysis was fully blinded.

the baryon density constraint is due to sensitivity to the shape of the matter power spectrum from baryon damping [74]. The constraint on  $\Omega_b$  from the CMB, by contrast, is also sensitive to the impact of baryons on the acoustic oscillations. This provides another avenue to test the predictions of our models from early Universe observations like the CMB with measurements from surveys like DES.

## OUTLOOK

The most precise constraints on dark energy properties require combining cosmological probes that include information from both geometry and growth across cosmic history. Thus far such diverse information was collected from different experiments, which were subject to different calibration and systematic errors. We have combined for the first time in DES the purely cosmographic SN and BAO measurements with the growth-sensitive weak lensing and galaxy clustering measurements to independently place strong constraints on the nature of dark energy. These results share a common set of calibration frameworks and blinding policy across probes. DES has independently constrained  $\Omega_m$ ,  $\Omega_b$ ,  $\Omega_\Lambda$ ,  $\sigma_8$ , and  $w$ , while allowing for a free neutrino mass. We have constrained  $w$  to a precision that is almost a factor of three better than the

previous best single-experiment result from the CMB, where the neutrino mass density was fixed [9]. We expect future DES results to lead to a further factor of 2-4 improvement in these constraints due to increased area, depth, and spectroscopic SNe followup in the final analyses, which will then be followed by subsequent order of magnitude advances from the major photometric surveys of the 2020s.

## ACKNOWLEDGEMENTS

Funding for the DES Projects has been provided by the DOE and NSF (USA), MEC/MICINN/MINECO (Spain), STFC (UK), HEFCE (UK), NCSA (UIUC), KICP (U. Chicago), CCAPP (Ohio State), MIFPA (Texas A&M), CNPQ, FAPERJ, FINEP (Brazil), DFG (Germany) and the Collaborating Institutions in the Dark Energy Survey.

The Collaborating Institutions are Argonne Lab, UC Santa Cruz, University of Cambridge, CIEMAT-Madrid, University of Chicago, University College London, DES-Brazil Consortium, University of Edinburgh, ETH Zürich, Fermilab, University of Illinois, ICE (IEEC-CSIC), IFAE Barcelona, Lawrence Berkeley Lab, LMU München and the associated Excellence Cluster Universe, University of Michigan, NOAO, University of Nottingham, Ohio State University, University of Pennsylvania, University of Portsmouth, SLAC National Lab, Stanford University, University of Sussex, Texas A&M University, and the OzDES Membership Consortium.

Based in part on observations at Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation.

The DES Data Management System is supported by the NSF under Grant Numbers AST-1138766 and AST-1536171. The DES participants from Spanish institutions are partially supported by MINECO under grants AYA2015-71825, ESP2015-66861, FPA2015-68048, SEV-2016-0588, SEV-2016-0597, and MDM-2015-0509, some of which include ERDF funds from the European Union. IFAE is partially funded by the CERCA program of the Generalitat de Catalunya. Research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Program (FP7/2007-2013) including ERC grant agreements 240672, 291329, and 306478. We acknowledge support from the Australian Research Council Centre of Excellence for All-sky Astrophysics (CAASTRO), through project number CE110001020, and the Brazilian Instituto Nacional de Ciência e Tecnologia (INCT) e-Universe (CNPq grant 465376/2014-2).

This manuscript has been authored by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the U.S. Department of Energy, Office of Science, Office of High Energy Physics. The United States Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-

exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

This research used resources of the National Energy Research Scientific Computing Center, a DOE Office of Science User Facility supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. This work also used resources on the CCAPP condo of the Ruby Cluster at the Ohio Supercomputing Center [75]. Plots in this manuscript were produced partly with MATPLOTLIB [76], and it has been prepared using NASA's Astrophysics Data System Bibliographic Services.

- 
- [1] A. G. Riess *et al.* (Supernova Search Team), *Astron. J.* **116**, 1009 (1998).
  - [2] S. Perlmutter *et al.* (Supernova Cosmology Project), *Astrophys. J.* **517**, 565 (1999).
  - [3] A. Albrecht *et al.*, ArXiv e-prints (2006), [arXiv:0609591](https://arxiv.org/abs/0609591).
  - [4] J. Frieman, M. Turner, and D. Huterer, *Ann. Rev. Astron. Astrophys.* **46**, 385 (2008).
  - [5] D. H. Weinberg, M. J. Mortonson, D. J. Eisenstein, C. Hirata, A. G. Riess, and E. Rozo, *Phys. Rep.* **530**, 87 (2013).
  - [6] D. Huterer and D. L. Shafer, *Rept. Prog. Phys.* **81**, 016901 (2018).
  - [7] W. Hu and S. Dodelson, *Ann. Rev. Astron. Astrophys.* **40**, 171 (2002).
  - [8] G. Hinshaw *et al.*, *ApJS* **208**, 19 (2013).
  - [9] Planck Collaboration *et al.*, ArXiv e-prints (2018), [arXiv:1807.06209](https://arxiv.org/abs/1807.06209).
  - [10] P. Astier *et al.*, *A&A* **572**, A80 (2014).
  - [11] L. Anderson *et al.*, *MNRAS* **441**, 24 (2014).
  - [12] S. Alam *et al.*, *MNRAS* **470**, 2617 (2017).
  - [13] S. W. Allen, A. E. Evrard, and A. B. Mantz, *ARA&A* **49**, 409 (2011).
  - [14] S. H. Suyu *et al.*, *MNRAS* **468**, 2590 (2017).
  - [15] R. Mandelbaum, ArXiv e-prints (2017), [arXiv:1710.03235](https://arxiv.org/abs/1710.03235).
  - [16] DES Collaboration *et al.*, *Phys. Rev. D* **98**, 043526 (2018).
  - [17] DES Collaboration *et al.*, ArXiv e-prints (2017), [arXiv:1712.06209](https://arxiv.org/abs/1712.06209).
  - [18] DES Collaboration *et al.*, ArXiv e-prints (2018), [arXiv:1811.02374](https://arxiv.org/abs/1811.02374).
  - [19] R. A. C. Croft and M. Dailey, ArXiv e-prints (2011), [arXiv:1112.3108](https://arxiv.org/abs/1112.3108).
  - [20] <http://hsc.mtk.nao.ac.jp/ssp/>.
  - [21] <http://kids.strw.leidenuniv.nl>.
  - [22] <http://www.darkenergysurvey.org>.
  - [23] <http://www.lsst.org>.
  - [24] <http://sci.esa.int/euclid>.
  - [25] <http://wfirst.gsfc.nasa.gov>.
  - [26] <http://www.sdss3.org/surveys/boss.php>.
  - [27] <https://www.sdss.org/surveys/eboss/>.
  - [28] <https://www.desi.lbl.gov>.
  - [29] <https://pfs.ipmu.jp>.
  - [30] B. Flaugher *et al.* (DES Collaboration), *Astron. J.* **150**, 150 (2015).
  - [31] S. Desai *et al.*, *ApJ* **757**, 83 (2012).
  - [32] I. Sevilla *et al.* (DES Collaboration), ArXiv e-prints (2011), [arXiv:1109.6741](https://arxiv.org/abs/1109.6741).
  - [33] J. J. Mohr *et al.*, in *Observatory Operations: Strategies, Pro-*

- cesses, and Systems II*, Proc. SPIE, Vol. 7016 (2008) p. 70160L.
- [34] E. Morganson *et al.* (DES Collaboration), *PASP* **130**, 074501 (2018).
- [35] A. Drlica-Wagner *et al.* (DES Collaboration), *ApJS* **235**, 33 (2018).
- [36] D. A. Goldstein *et al.* (DES Collaboration), *AJ* **150**, 82 (2015).
- [37] R. Kessler *et al.* (DES Collaboration), *AJ* **150**, 172 (2015).
- [38] E. Huff and R. Mandelbaum, ArXiv e-prints (2017), [arXiv:1702.02600](https://arxiv.org/abs/1702.02600).
- [39] E. S. Sheldon and E. M. Huff, *Astrophys. J.* **841**, 24 (2017).
- [40] J. Zuntz *et al.* (DES Collaboration), *MNRAS* **481**, 1149 (2018).
- [41] N. Benítez, *Astrophys. J.* **536**, 571 (2000).
- [42] B. Hoyle *et al.* (DES Collaboration), *MNRAS* **478**, 592 (2018).
- [43] E. Rozo *et al.* (DES Collaboration), *MNRAS* **461**, 1431 (2016).
- [44] J. Elvin-Poole *et al.* (DES Collaboration), *Phys. Rev. D* **98**, 042006 (2018).
- [45] E. Krause and T. Eifler, *MNRAS* **470**, 2100 (2017).
- [46] E. Krause *et al.* (DES Collaboration), ArXiv e-prints (2017), [arXiv:1706.09359](https://arxiv.org/abs/1706.09359).
- [47] N. MacCrann *et al.* (DES Collaboration), *MNRAS* **480**, 4614 (2018).
- [48] J. Prat *et al.* (DES Collaboration), *Phys. Rev. D* **98**, 042005 (2018).
- [49] M. A. Troxel *et al.* (DES Collaboration), *Phys. Rev. D* **98**, 043528 (2018).
- [50] J. Zuntz *et al.*, *Astronomy and Computing* **12**, 45 (2015).
- [51] C. D’Andrea *et al.* (DES Collaboration), in prep. (2018).
- [52] D. Brout *et al.*, ArXiv e-prints (2018), [arXiv:1811.02378](https://arxiv.org/abs/1811.02378) [astro-ph.IM].
- [53] D. L. Burke *et al.*, *AJ* **155**, 41 (2018).
- [54] J. Lasker *et al.*, ArXiv e-prints (2018), [arXiv:1811.02380](https://arxiv.org/abs/1811.02380).
- [55] R. Kessler *et al.*, ArXiv e-prints (2018), [arXiv:1811.02379](https://arxiv.org/abs/1811.02379).
- [56] D. Scolnic and R. Kessler, *ApJ* **822**, L35 (2016).
- [57] D. Brout *et al.*, ArXiv e-prints (2018), [arXiv:1811.02377](https://arxiv.org/abs/1811.02377).
- [58] E. Macauley *et al.*, ArXiv e-prints (2018), [arXiv:1811.02376](https://arxiv.org/abs/1811.02376).
- [59] M. Hicken *et al.*, *ApJ* **700**, 1097 (2009).
- [60] M. Hicken *et al.*, *ApJS* **200**, 12 (2012).
- [61] C. Contreras *et al.*, *AJ* **139**, 519 (2010).
- [62] D. M. Scolnic *et al.*, *ApJ* **859**, 101 (2018).
- [63] M. Crocce *et al.* (DES Collaboration), *MNRAS*, 2411 (2018).
- [64] S. Avila *et al.* (DES Collaboration), *MNRAS* **479**, 94 (2018).
- [65] A. J. Ross *et al.* (DES Collaboration), *MNRAS* **472**, 4456 (2017).
- [66] K. C. Chan *et al.* (DES Collaboration), *MNRAS* **480**, 3031 (2018).
- [67] H. Camacho *et al.* (DES Collaboration), ArXiv e-prints (2018), [arXiv:1807.10163](https://arxiv.org/abs/1807.10163).
- [68] Planck Collaboration, P. A. R. Ade, *et al.*, *A&A* **594**, A13 (2016).
- [69] Planck Collaboration, P. A. R. Ade, *et al.*, *A&A* **594**, A15 (2016).
- [70] Planck Collaboration, N. Aghanim, *et al.*, *A&A* **594**, A11 (2016).
- [71] D. Scolnic *et al.*, *ApJ* **815**, 117 (2015).
- [72] F. Beutler *et al.*, *Mon. Not. Roy. Astron. Soc.* **416**, 3017 (2011).
- [73] A. J. Ross *et al.*, *Mon. Not. Roy. Astron. Soc.* **449**, 835 (2015).
- [74] This constraint on  $\Omega_b$  is not prior driven, despite being degenerate with  $A_s$ , and  $n_s$ , which is not constrained.
- [75] OSC, “Ohio supercomputer center,” <http://osc.edu/ark:/19495/f5s1ph73> (1987).
- [76] J. D. Hunter, *Computing In Science & Engineering* **9**, 90 (2007).